



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# The impact of an observationally based surface emissivity dataset on the simulation of Microwave Sounding Unit Temperatures

J. J. Hnilo, L. Litten , B. D. Santer, J. R. Christy

August 15, 2005

86th American Meteorological Society Annual Meeting  
Atlanta, GA, United States  
January 29, 2006 through February 2, 2006

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

## The impact of an observationally based surface emissivity dataset on the simulation of Microwave Sounding Unit Temperatures.

J. J. Hnilo, L. Litten, B. D. Santer and J. R. Christy

Relatively few studies have attempted to simulate synthetic MSU temperatures with use of a radiation model. Most employ the simpler and computationally less-expensive method of applying a static, global-mean weighting function to three-dimensional profiles of atmospheric temperature. Both approaches require a number of key assumptions. One of the major assumptions relates to surface emissivity. To date, two different strategies have been used for prescribing surface emissivity values. The first assumes a fixed global surface emissivity, while the second specifies separate (time-invariant) emissivity values for land and ocean.

In this research, we introduce space- and time-dependence to the specified emissivity fields, using recent observationally-based estimates of surface emissivity changes over 1988 to 2000. We use a radiative transfer code to explore the impact of this more complex treatment of surface emissivity. This sensitivity analysis is performed with monthly-mean fields of surface temperature, atmospheric temperature, and moisture taken from multiple reanalyses. Our goal is to quantify the possible impact of emissivity changes on global-scale estimates of tropospheric temperature trends (*e.g.*, trends estimated from MSU channel 2 and MSU 2LT), and to document the sensitivity of synthetic MSU temperatures to a variety of input data and processing choices.

This work is supported under the auspices of the Office of Science, U.S. Department of Energy at the University of California Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48